



FAVE Sección Ciencias Veterinarias
ISSN: 1666-938X
ISSN: 2362-5589
favecv@gmail.com
Universidad Nacional del Litoral
Argentina

Descriptive study of congenital anomalies encountered in ruminants in Elazig region of Turkey

Polat, E.

Descriptive study of congenital anomalies encountered in ruminants in Elazig region of Turkey

FAVE Sección Ciencias Veterinarias, vol. 21, 2022

Universidad Nacional del Litoral, Argentina

Disponible en: <https://www.redalyc.org/articulo.oa?id=617770132001>

DOI: <https://doi.org/10.14409/favecv.2022.1.e0001>



Esta obra está bajo una Licencia Creative Commons Atribución-NoComercial-CompartirIgual 4.0 Internacional.

Descriptive study of congenital anomalies encountered in ruminants in Elazig region of Turkey

Estudio descriptivo de anomalías congénitas encontradas en rumiantes en la región de Elazig en Turquía

Polat, E.
Firat University, Turquía
erenpolat@firat.edu.tr

DOI: <https://doi.org/10.14409/favecv.2022.1.e0001>
Redalyc: <https://www.redalyc.org/articulo.oa?id=617770132001>

Recepción: 14 Junio 2021
Aprobación: 28 Febrero 2022

ABSTRACT:

Congenital malformations are abnormalities of structure, formation or function observable at birth that causes great economic losses in livestock industry. In this study, we describe the incidence of the main congenital anomalies encountered in the last year (June 2020-June 2021) in ruminants from Elazig region according to species, breed, age and sex, number of pregnancies, rectal examination, and method of conception employed. A total of 79 congenital anomalies encountered in 73 animals during this study were evaluated. Of 73 animals most of congenital anomalies were calves (84.93%) and males (68.49%). It was determined that the most common type of congenital anomaly in calves was hernia umbilicalis (21.51%) while in lambs and goat kids was atresia ani. Most of these anomalies were formed in the abdominal wall and organs of gastrointestinal system (46.84%). Congenital anomalies in calves and lambs were more frequently when they came from the first pregnancy of their dams, while in goat kids when their dam had the second pregnancy. Most of small ruminant that had congenital anomalies came from dams with multiple gestation. Rectal examinations performed in cattle during the early stages of gestation was probably one of a predisposing factor for the formation of congenital malformations in this specie. More studies that include the participation of diagnostic laboratories are needed to improve the characterization of congenital anomalies that contribute to the implementation of preventive and control measures in the herds of these species in our region.

KEYWORDS: anomaly, calf, congenital, goat kids, lamb, ruminant.

RESUMEN:

Las malformaciones congénitas son anomalías de estructura, formación o función observables al nacer que provocan grandes pérdidas económicas en la industria ganadera. En este estudio, se investigó la incidencia de las principales anomalías congénitas encontradas en el último año (junio de 2020-junio de 2021) en rumiantes en la región de Elazig, según especie, raza, edad, sexo, número de preñeces, examen rectal y método de concepción empleado. Se evaluaron un total de 79 anomalías congénitas encontradas en 73 animales. De 73 animales con anomalías congénitas, la mayoría eran terneros (84,93%) y machos (68,49%). Se determinó que el tipo de anomalía más común en los terneros fue la hernia umbilical (21,51%) mientras que en corderos y cabritos fue la atresia anal. La mayoría de estas anomalías se formaban principalmente en los órganos de la pared abdominal y órganos del sistema gastrointestinal (46,84%). Las anomalías congénitas en terneros y corderos eran más frecuentes cuando provenían de las madres de primera preñez, mientras que en los cabritos cuando las madres tenían su segunda preñez. La mayoría de los rumiantes menores con anomalías congénitas provenían de madres con gestaciones múltiples. El examen rectal realizado en las vacas durante los primeros estadios de la gestación, fue probablemente uno de los factores predisponentes para la formación de malformaciones congénitas en esta especie. Se requieren más estudios que incluyan la participación de laboratorios de diagnóstico para mejorar la caracterización de las anomalías congénitas y contribuir a la implementación de medidas de prevención y control en los rodeos de estas especies en nuestra región.

PALABRAS CLAVE: anomalía, ternero, congénito, cabrito, cordero, rumiante.

NOTAS DE AUTOR

Eren POLAT, Firat Universitesi Veteriner Fakültesi, 23119 Elazig - Turkey

INTRODUCTION

Congenital anomalies are structural and functional disorders that occur in newborns under the influence of genetic and environmental factors during pregnancy (Aksoy et al. 2006; Belge et al. 2000; Demiraslan et al. 2014; Kaya et al. 2011; Niwas et al. 2020; Tasal and Aytekin 2015). It is very difficult to determine the incidence and etiology of congenital anomalies because animals with butchery value are mostly not treated. Congenital anomalies may be caused by genetic factors such as chromosomal defects and mutations, as well as by environmental factors such as feed and water taken during pregnancy, intrauterine infections and iatrogenic factors like rectal examinations in the early period of pregnancy (Aksoy et al. 2006; Albarella et al. 2017; Bademkiran et al. 2009; Belge et al. 2000; Chauhan et al. 2011; Durmus and Polat 2019; Günay et al. 2020; Hussein 2010; Kılıç et al. 2016; Niwas et al. 2020; Ogurtan et al. 1997; Ozaydın et al. 2006; Syed and Shanks 1992; Tasal and Aytekin 2015; Varol et al. 2018; Yayla et al. 2016; Yayla et al. 2017) or drug administration. Although congenital anomalies mostly occur in the digestive system, and musculoskeletal system, it also occurs less frequently in the urogenital system and skin. Atresia ani, bouleture and hernia umbilicalis are among the most common congenital anomalies (Aksoy et al. 2006; Belge et al. 2000; Fubini 2004; Kaya et al. 2011; Lotfi and Shahryar 2009).

The aim of this study was to describe the main congenital anomalies encountered in ruminants in Elazig region of Turkey.

MATERIAL AND METHODS

Ethical Statement and Animal Material

This study was conducted after the approval of Firat University Animal Experiments Local Ethics Committee in accordance with ethical principles (Meeting no: 2020/8, date 10.06.2020). The study was performed in the ruminant livestock business (cattle, sheep, goats etc.) in Elazig region of Turkey. In this study a total of 79 congenital anomalies encountered in 73 animals were evaluated.

Evaluation of Congenital Anomalies

A total of 73 animals with congenital anomalies were examined macroscopically and an exhaustive description of them were performed. In each case, the specie, breed, sex, age and the origin were registered. In addition, a detailed and comprehensive questionnaire was performed in order to obtain more information for this study (Table 1).

TABLE 1.
Questions asked to breeders to evaluate the etiology of congenital anomalies

	Questions
1	Which method was used to obtain pregnancy? O Artificial insemination O Natural fertilization
2	How many times did the animal give birth before that?
3	Has a rectal examination been performed during pregnancy? If so, what period of pregnancy was it? O Rectal examination was not performed during pregnancy O Rectal examination was performed in the first trimester of pregnancy. O Rectal examination was performed in the second trimester of pregnancy. O Rectal examination was performed in the last trimester of pregnancy
4	Did any illness occur during pregnancy?
5	Were animals fed in a controlled manner during pregnancy? O Yes (Animals were fed in a controlled manner in barns and pens) O No (Animals were fed with pasture grasses)
6	Have twins or more offspring born at birth? If so, is there any abnormality was observed in the offspring?

The rates of congenital anomalies according to the species, breed, sex age and the organ/system affected were determined.

RESULTS

Between June 2020 and June 2021, descriptive study of congenital anomalies encountered in ruminant farms in Elazig region were performed. During the study, a total of 79 congenital anomalies were detected in 73 animals (Fig. 1- a,b,c,d). Congenital anomalies were detected in 84.93% (n= 62) calves, 9.59% (n= 7) in lambs, and 5.48% (n= 4) in goat kids. These anomalies were detected in 68.49% (n= 50) male animals and 31.51% (n= 23) in female animals. The distribution of congenital anomalies according to species, breed and gender are presented in *table 2* and *table 3*.




FIGURE 1.
Some congenital anomalies: atresia ani (a) anophthalmia (b) mandibular malformation brachygnathia inferior (c) cheliognatopalatoschisis (d)

TABLE 2.
Distribution of congenital anomalies according to the species and breeds of animals

Animal Species	Anomaly Cases		Animal Breeds	Anomaly Cases	
	n	Frequency		n	Frequency
Calves	62	84.93 %	Simmental	51	82.26 %
			Holstein	11	17.74 %
Lambs	7	9.59 %	White Karaman	6	85.71 %
			Red Karaman	1	14.29 %
Goat kids	4	5.48 %	Hair Goat	3	75 %
			Aleppo Goat	1	25 %
Total	73	100 %	Total	73	100%

TABLE 3.
Distribution of congenital anomalies by gender of animals

Animal Species	Gender of Animals			
	Male		Female	
	n	Frequency	n	Frequency
Calves	43	69.35 %	19	30.65 %
Lambs	4	57.14 %	3	42.86 %
Goat kids	3	75 %	1	25 %
Total	50	68.49 %	23	31.51 %

In this study, hernia umbilicalis (21.51%), atresia ani (15.18%) and bouleture (flexor tendon deformity of the metacarpophalangeal joint) (12.65%) were f the most common congenital anomalies found . The systems that most frequently included congenital anomalies were the abdominal wall-gastrointestinal system

(46.84%) and the musculoskeletal system (30.38%). The distribution of congenital anomalies according to their types and systems involved is presented in *Table 4*.

TABLE 4.
Distribution of congenital anomalies according to their types and systems involved

Anomaly Types	Distribution of Anomalies		Systems with Anomalies	Distribution of Anomalies	
	n	Frequency		n	Frequency
Hamartoma	1	1.27 %	Formation Anomalies in the Head Region	10	12.65 %
Mandibular malformation	3	3.80 %			
Cheliognatopalatoschisis	2	2.53 %			
Anophthalmia	1	1.27 %			
Microphthalmia	1	1.27 %			
Cyst Dermoid	2	2.53 %			
Atresia ani	12	15.18 %	Abdominal Wall and Gastrointestinal organs	37	46.84 %
Hernia umbilicalis	17	21.51 %			
Atresia coli	6	7.59 %			
Schistosoma reflexum	2	2.53 %			
Coccygeal agenesis	7	8.86 %	Musculoskeletal System	24	30.38 %
Arqure (flexor tendon deformity in the carpal joint)	3	3.80 %			
Bouleture (flexor tendon deformity of the metacarpophalangeal joint)	10	12.65 %			
Tarsal hyperextension	3	3.80 %			
Arthrogryposis	1	1.27 %			
Urethral Dilatation	1	1.27 %			
Tracheal Collapse	2	2.53 %	Respiratory System	2	2.53 %
Hydrocephalus	1	1.27 %	Nervous System	1	1.27 %
Rectourethral fistula	1	1.27 %	Multiple System	4	5.06 %
Rectovaginal fistula	3	3.80 %			
Total	79	100 %	Total	79	100 %

Most of congenital anomalies were detected between 0-7 days old (73.97%), and secondarily when the animals were more than 14 days old (17.81%). The relationship between the age and the presence of congenital anomalies in each animal species is presented in *Table 5*.

TABLE 5.
Relationship between the age and congenital anomalies in each animal species

Age	Calves		Lambs		Goat kids		Total	
	n	Frequency	n	Frequency	n	Frequency	n	Frequency
0-7 days	44	70.97 %	6	85.71 %	4	100 %	54	73.97 %
7-14 days	6	9.68 %	-	-	-	-	6	8.22 %
> 14 days	12	19.35 %	1	14.29 %	-	-	13	17.81 %
Total	62	100 %	7	100 %	4	100 %	73	100 %

A total of 66 congenital anomalies were found in 62 calves. Most of them consisted of hernia umbilicalis (25.75%), bouleture (13.63%), coccygeal agenesis (10.60%) and atresia coli (9.09%). The distribution of congenital anomalies encountered in calves is presented in *table 6*.

TABLE 6.
Distribution of congenital anomalies in 66 calves according to their types

Anomaly Types	Distribution of Anomalies	
	n	Frequency
Hamartoma	1	1.52 %
Mandibular malformation	3	4.54 %
Cheliognatopalatoschisis	2	3.03 %
Anophthalmia	1	1.52 %
Microphthalmia	1	1.52 %
Cyst Dermoid	2	3.03 %
Atresia ani	3	4.54 %
Hernia umbilicalis	17	25.75 %
Atresia coli	6	9.09 %
Schistosoma reflexum	2	3.03 %
Coccygeal agenesis	7	10.60 %
Argure	3	4.54 %
Bouleture	9	13.63 %
Tarsal hyperextension	2	3.03 %
Arthrogryposis	1	1.52 %
Urethral Dilatation	1	1.52 %
Tracheal Collapse	2	3.03 %
Hydrocephalus	1	1.52 %
Rectourethral fistula	1	1.52 %
Rectovaginal fistula	1	1.52 %
Total	66	100 %

Atresia ani was the most common congenital anomaly in lambs (66.67%) and goat kids (75%). (Fig. 2- a,b).




FIGURE 2. Percentage of congenital anomalies encountered in lambs a and goat kids b

It was determined that 53.23% of cattle that had offspring with congenital anomalies were pregnant by artificial insemination. in contrast to sheep and goats who were pregnant by natural breeding. The relationship between the congenital anomalies of the offspring and the way of conception received in each animals species is presented in *table 7*.

TABLE 7. Relationship and distribution between the offspring with congenital anomalies and the way of conception in each animal species

Animal Species	Artificial insemination		Breeding	
	n	Frequency	n	Frequency
Calve	33	53.23 %	29	46.77 %
Lamb	-	-	7	100 %
Goat kids	-	-	4	100 %
Total	33	45.21 %	40	54.79 %

It was determined that the presence of offspring with congenital anomalies in cattle, sheep and goats in their first pregnancy was 56.45%, 14.29% and 75 %, in each case, respectively... It was found that 53.42% of all animals species had offspring with congenital anomalies in their first pregnancy . The relationship between the number of pregnancies and the offspring with congenital anomalies is presented in *table 8*.

TABLE 8.

Relationship and frequency between number of pregnancies and offspring with congenital anomalies

Number of pregnancies	Cattle		Sheep		Goat		Total	
	n	Frequency	n	Frequency	n	Frequency	n	Frequency
First	35	56.45 %	1	14.29 %	3	75 %	39	53.42 %
2-3	18	29.03 %	5	71.42 %	-	-	23	31.51 %
≥ 4	9	14.52 %	1	14.29 %	1	25 %	11	15.07 %

It was determined that 61.29% of cattle that had offspring with congenital anomalies received rectal examination for the diagnosis of pregnancy in the early period of gestation. The relationship between the period of rectal examinations performed during pregnancy and the birth of offspring with congenital anomalies is presented in *table 9*.

TABLE 9.

The relationship of rectal examination periods during pregnancy with birth of offspring with congenital anomalies

Rectal Examination performed during gestation period	Cattle		Sheep		Goat		Total	
	n	Frequency	n	Frequency	n	Frequency	n	Frequency
RM (-)	11	17.74 %	7	100 %	4	100 %	22	30.14 %
Early period	38	61.29 %	-	-	-	-	38	52.06 %
Middle period	7	11.29 %	-	-	-	-	7	9.59 %
Last period	6	9.68 %	-	-	-	-	6	8.21 %

It was found that 83.56% of the animals that had offspring with congenital anomalies were grazing on pasture. It was determined that 82.19% of the animals did not have any disease during pregnancy.

It was found that 19.18% of the animals with congenital anomalies were born as a result of twin (cattle) or multiple pregnancy (small ruminants). In small ruminants, the percentage of congenital anomalies was higher in lamb (85.71%) than goatkids (75%) (*Table 10*).

TABLE 10.

The relationship between birth of offspring with congenital anomalies and the situations of giving birth to a single or twin multiple birth

Animal Species	Twin/multiple offspring		Single offspring	
	n	Frequency	n	Frequency
Calve	5	8.06 %	57	91.94 %
Lamb	6	85.71 %	1	14.29 %
Goat kids	3	75 %	1	25 %
Total	14	19.18 %	59	80.82 %

DISCUSSION

Congenital anomalies are structural and functional pathologies related with genetic origin and/or environmental factors that acts during the pregnancy (Aksoy et al. 2006; Belge et al. 2000; Demiraslan et al. 2014; Kaya et al. 2011; Niwas et al. 2020; Tasal and Aytekin 2015). The incidence of congenital anomalies in ruminants ranges from 26.8% to 40.48% (Fubini 2004; Kaya et al. 2011; Ogurtan et al. 1997). Some congenital anomalies can cause death, while others can only create a structural defect without affecting

vital functions (Kaya et al. 2011). In this study, the aim was to describe the main congenital anomalies encountered in ruminants from the Elazig region in Turkey.

Most of the congenital anomalies encountered in ruminants are seen in calves (Aksoy et al. 2006; Doğan and Sındak 2013). According to this, Aksoy et al. (2006), reported that 83.3% of congenital anomalies encountered in ruminants in Kars region are seen in calves. In addition, this author (Aksoy et al. 2006), found that the rate of congenital anomalies was 14.8% in lambs and 1.9% in goat kids. Doğan and Sındak (2013), reported that 72.5% of congenital anomalies encountered in ruminants in Nizip region are seen in calves, being the rate of congenital anomalies of 7.5% in lambs and 20% in goat kids, respectively. In this study, it was found that 84.93% of congenital anomalies were seen in calves 9.59% in lambs and 5.48% in goat kids. The higher rate of congenital anomalies described by Doğan and Sındak (2013), in goats compared to other studies is related to the fact that the breeding of this species is higher in Nizip region compared to Elazig and Kars regions.

Some researchers reported that congenital anomalies encountered in calves are more common in Swiss Brown breed (Karabulut et al. 2001; Kaya et al. 2011). Some researchers reported that congenital anomalies are more common in Simmental calves.¹ Kaya et al. (2011), reported that 47.42% of the congenital anomalies encountered in calves in Erzurum region occurred in the Swiss Brown breed. On the other hand, Karabulut et al. (2001), reported that 58.94% of the congenital anomalies encountered in calves in Elazig region occurred in the Swiss Brown breed. Aksoy et al. (2006), reported that 32.1% of congenital anomalies are in Simmental calves. Belge et al. (2000), reported that congenital anomalies are more common in crossbred calves. In this study, it was determined that 82.26% of the congenital anomalies encountered in calves occurred in the Simmental breed. All these studies suggest that congenital anomalies have not a breed predisposition. The reason for the difference between these studies is thought to be related with the populations of cattle breeds according to the regions.

Previous studies demonstrated that the incidence of congenital anomalies is higher in male animals species (Aksoy et al. 2006; Belge et al. 2000; Kaya et al. 2011). Aksoy et al. (2006), in their study on ruminants in Kars region, reported that 67.55% of the animals with congenital anomalies were male, 64.67% were calves, 76.92% lambs, and 100% goat kids. Kaya et al. (2011), reported that 56.70% of the animals with congenital anomalies were male in their study on calves in Erzurum region. In a study conducted in the Van region, Belge et al. (2000), reported that 59.60% of the animals with congenital anomalies were males. In this study, 69.35% of calves, 57.14% of lambs and 75% of goat kids were male.

Some researchers reported higher incidence of congenital anomalies in the musculoskeletal system (Kaya et al. 2011; Ogurtan et al. 1997; Ozaydın et al. 2006; Syed and Shanks 1992). In agreement with other authors (Aksoy et al. 2006, Dogan and Sındak 2013) in this study the prevalence of congenital anomalies in the abdominal wall and gastrointestinal system (46.84%) was higher than other types of anomalies. The rate of musculoskeletal anomalies was 30.38% being the second system with congenital anomalies.

It has been reported that the most common congenital anomalies in calves are hernia umbilicalis, atresia ani and bouleture (Aksoy et al. 2006; Belge et al. 2000; Fubini 2004; Kaya et al. 2011; Lotfi and Shahryar 2009). However, Karabulut et al. (2001), reported that the most common anomalies in calves in Elazig region were bouleture (45.26%), cyst dermoid (17.89%) and arqure (17.89%). Doğan and Sındak (2013), reported that the most common congenital anomalies in calves in the Nizip region were hernia umbilicalis and cyst dermoid. In this study, the most common were hernia umbilicalis (25.76%), bouleture (13.63%) and coccygeal agenesis (10.60%). The rate of cyst dermoid cases was 3.03%.

Some researchers reported that the most common congenital anomalies in lambs are atresia ani, and in goat kids the urogenital system, respectively. (Aksoy et al. 2006; Doğan and Sındak 2013; Ogurtan et al. 1997). In this study, it was determined that atresia ani was the most common cases of congenital anomaly in lambs (66.7%) and goat kids (75%) however, urogenital system anomalies were not observed in goat kids.

In some studies (Aksoy et al. 2006; Bahr et al. 2003; Belge et al. 2000; Fırat et al. 2005; Harper et al. 1990; Miura et al. 1990; Ogurtan et al. 1997; Stanley 1993; Tsuda et al. 2004), it has been reported that the incidence of congenital anomalies is lower in the offspring of animals conceived by artificial insemination. In this study, it was determined that 53.23% of the calves with congenital anomalies became pregnant by artificial insemination being the incidence of congenital anomalies lower in offspring conceived by natural mating than in offspring conceived by artificial insemination.

Rectal examinations for pregnancy diagnosis performed in the early stages of gestation can be the cause of many congenital anomalies, especially intestinal atresia (Aksoy et al. 2006; Atalan et al. 2003; Belge et al. 2000; Demiraslan et al. 2014; Hendrickson et al. 1992; Kaya et al. 2011; Longeri et al. 2003; Newman et al. 1999). In this study, rectal examination was performed in the early period of gestation in 61.29% of cattle that had offspring with congenital anomalies, but atresi ani was not registered in these calves. Probably this practice was one of the predisposing factor for congenital malformations.

Congenital anomalies in calves and lambs were more frequently when they came from the first pregnancy of their dams, and in goat kids that they came from the second dam pregnancy. Most of small ruminant that had congenital anomalies came from dams with multiple gestation. frequently

Congenital anomalies are pathologies frequently present in ruminant breeding herds all over the world. Although the prevalence of congenital anomalies in ruminants varies according to the regions, it had been recorded between 26.8-40.48%. This study, describe the main congenital anomalies encountered in ruminants in the Elazığ region of Turkey and the relationship between species, race, sex, age, number of pregnancies, rectal examination, and method of conception employed . More studies that include the participation of diagnostic laboratories are needed to improve the characterization of congenital anomalies that could contribute to the implementation of preventive and control measures

REFERENCES

- Aksoy O, Kılıç E, Öztürk S, Özyaydın I, Kurt B, Baran V. 2006. Congenital Anomalies Encountered in Calves, Lambs and Kids: 1996-2005 (262 Cases). *Kafkas Univ. Vet. Fak. Derg.* 12: 147-154.
- Albarella S, Ciotola F, D'Anza E, Coletta A, Zicarelli L, Peretti V. 2017. Congenital Malformations in River Buffalo. *Animals* 7: 1-15.
- Atalan G, Özyaydın I, Kılıç E, Cihan M, Kamiloğlu A. 2003. Occurrence of intestinal atresia and their operative treatments in calves: 54 cases (1992-2000). *Kafkas Univ. Vet. Fak. Derg.* 9: 113-118.
- Bademkiran S, İçen H, Kurt D. 2009. Congenital rectovaginal fistula with atresia ani in a heifer. *Y.Y.Ü. Vet. Fak. Derg.* 20: 61-64.
- Bahr C, Kuiper H, Distl O. 2003. Bilateral anophthalmia associated with further anomalies of the head in German Holstein calves. *Dtsch. Tierärztl. Wochenschr.* 110: 454-456.
- Belge A, Gönenci R, Biricik HS, Ormancı S. 2000. Buzağlarda doğumsal anomali olguları. *Y.Y.Ü. Vet. Fak. Derg.* 11: 23-26.
- Chauhan PM, Parmar VR, Patel TP, Thakor KB, Parikh SS. 2011. Atresia ani: A congenital defect and its successful management in non-descript calf. *IJAVMS* 5: 520-522.
- Demiraslan Y, Aslan K, Gürbüz I, Özen H. 2014. Multiple Congenital Anomalies in a Simmental Calf. *Kafkas Univ. Vet. Fak. Derg.* 20: 629-632.
- Dogan H, Sindak N. 2013. The incidence of anomalies in calves, lambs and goat kids in the district of Nizip and its villages and determination of some biochemical parameters in these cases. *Harran Üniv. Vet. Fak. Derg.* 2: 61-66.
- Durmus AS, Polat E. 2019. A Case of Atresia Ani and Rectourethral Fistula in a Simmental Calf. *Harran Üniv. Vet. Fak. Derg.* 8: 246-248.
- Elias E, Bennett R. 1992. Congenital defects in Awassi fat-tailed lambs. *Small Rum. Res.* 8 (1-2): 141-150.

- Fırat I, Yıldız F, Özsoy S. 2005. A case of congenital flexure of fetlock in a Holstein calf. *İ.Ü. Vet. Fak. Derg.* 31: 103-108.
- Fubini SL. 2004. Surgery of the Perineum. In: *Farm Animal Surgery*, 1st ed., Philadelphia, USA: Saunders Publishing. Pp. 399-403.
- Günay C, Polat E, Sağlıyan A. 2019. Clinical, Radiological, Ultrasonographic Diagnosis and Treatment of Urethral Dilatation Case in a Simental Calf. *Harran Üniv. Vet. Fak. Derg.* 9: 082-085, 2019.
- Harper P, Plant JW, Unger DB. 1990. Congenital biliary atresia and jaundice in lambs and calves. *Aust Vet. J.* 67: 18-22.
- Hendrickson DA, Rackstraw PC, Ducharme NG. 1992. Surgical repair of atresia jejuni in two calves. *JAVMA* 201: 594-596, 1992.
- Hussein RMN. 2010. Congenital anomalies in cattle and buffalo within mudaina city in Basrah province between period 2007-2009. *Kufa J. Vet. Sci.* 1: 207-218.
- Karabulut E, Unsaldi S, Durgun T. 2001. 1991-2000 yılları arasında Fırat Üniversitesi Veteriner Fakültesi Cerrahi Kliniği'ne getirilen buzağlardaki doğumsal anomali olguları. *F.Ü. Sağ. Bil. Derg.* 15: 367-374.
- Kaya M, Okumus Z, Dogan E, Cetin EM, Yanmaz LE. 2011. Prevalence and Survival Rate of Congenital Malformations in Calves in Erzurum Districts. *F.Ü. Sağ. Bil. Vet. Derg.* 25: 83-93.
- Kılıç E, Özaydın I, Aksoy O, Yayla S, Sözmen M. 2006. Multiple urogenital system anomalies in three calves. *Kafkas Üniv. Vet. Fak. Derg.* 12: 193-197.
- Longeri M, Perrone T, Bongioni G, Bona M, Zanotti M, Galli A. 2003. Survival motor neuron (SMN) polymorphism in relation to congenital arthrogyriposis in two Piedmont calves. *Genet. Sol. Evol.* 35: 167-175.
- Lotfi A, Shahryar HA. 2009. The case report of taillessness in Iranian female calf (A congenital abnormality). *AJAVA* 4: 47-51.
- Miura R, Kubo M, Goto Y, Kono Y. 1990. Hydranencephaly-cerebellar hypoplasia in newborn calf after infection of its dam with Chuzan virus. *Nippon Juigaku Zasshi.* 52: 689-694.
- Newman SJ, Bailey TL, Jones JC, DiGrassie WA, Whittier WD. 1999. Multiple congenital anomalies in calf. *J. Vet. Diagn. Invest.* 11: 368-371.
- Niwas R, Najjar R, Priyanka, Dinesh, Saharan S. 2020. Common congenital surgical affections in large ruminants. *The Pharma Innov. Journ.* 9: 484-490.
- Oğurtan Z, Alkan F, Koç Y. 1997. Ruminantlarda konjenital anomaliler. *Türk Vet. Derg.* 9: 24-28.
- Özaydın I, Kılıç E, Aksoy O, Cihan M, Güngör E. 2006. Bir buzağda üçlü malformasyon: atrezi ani, rektouretral fistül ve pygomelia. *Kafkas Üniv. Vet. Fak. Derg.* 12: 189-191.
- Stanley DM. 1993. Congenital abnormalities. *Vet. Clin. North Am.: Food Anim. Pract.* 9: 1-206.
- Syed M, Shanks RD. 1992. Atresia coli inherited in holstein cattle. *J. Dairy Sci.* 75: 1105-1111.
- Tasal I, AYTEKİN O. 2015. A case of brachygnathia inferior (micrognathia) in a calf. *Van. Vet. J.* 26: 157-159.
- Tsuda T, Yosida K, Ohashi S, Yanase T, Sueyoshi M, Kamimura S, Misumi K, Hamana K, Sakamoto H, Yamakawa M. 2004. Arthrogyriposis, hydranencephaly and cerebellar hypoplasia syndrome in neonatal calves resulting from intrauterine infection with Anio virus. *Vet. Res.* 35: 531-538.
- Varol K, Atalan G, Günes V, Alpman U, Yönez MK. 2018. A case of atresia ani in an anatolian water buffalo calf. *Erciyes Üniv. Vet. Fak. Derg.* 15: 271-275.
- Yayla S, Kılıç E, Baran V, Ozen H. 2016. A case of congenital rectourethral fistula, urethral dilatation and segmental urethral agenesis encountered in a calf. *Harran Üniv. Vet. Fak. Derg.* 5: 70-73.
- Yayla S, Kılıç E, Ozen H, Baran V, Aydın U, Karakurt E. 2017. Simental ırkı bir buzağda çoklu ürogenital sistem anomalisi. *Harran Üniv. Vet. Fak. Derg.* 6: 95-98.

NOTES

Financial SupportNo financial support was received from any person or institution in this study.

Conflict of InterestThe authors declared that there is no conflict of interest.

Author ContributionsEP did all of the planning, execution and writing of the work.