

REFERENCIAS BIBLIOGRÁFICAS

- Abee, T. (1995). Pore-forming bacteriocins of Gram-positive bacteria and self-protection mechanisms of producer organisms. FEMS Microbiol. Lett. 129, 1-10.
- Amsterdam, D. (1996). Susceptibility testing of antimicrobials in liquid media. In: Loman, V.,(Ed). Antibiotics in laboratory medicine, 4th Ed. Williams and Wilkins, Baltimore, Md. p:52-111.
- Andersen, E.L; Diep, D.B; Nes, I.F; Eijsink, V.G.H; Nissen-Meyer, J. (1998). Antagonistic activity of *Lactobacillus plantarum* C11: two new two-peptide bacteriocins, Plantaricins EF and JK, and the induction factor Plantaricin A - Appl. Environ. Microbiol. 64 (6): 2269-2272.
- Atrih, A; Rekhif, N; Moir, A.J; Lebrihi, A.; Lefebvre, G. (2001). Mode of action, purification and aminoacid sequence of Plantaricin C19, an anti-Listeria bacteriocin produced by *Lactobacillus plantarum* C19 - Int. J. Food Microbiol. 68 (1)-2:93-104.
- Berry, E.D; Liewen, M.B; Mandigo, R.W; and Hutkins R.W. (1990). Inhibition of *Listeria monocytogenes* by bacteriocin-producing *Pediococcus* during the manufacture of fermented semidry sausages - J. Food. Prot . 52:194-197.
- Beukes, M; Bierbaum, G; Sahl, H.G and Hastings, J.W. (2000). Purification and partial characterization of a murein hydrolase, Millericin B, produced by *Streptococcus milleri* NMSCC061- Appl Environ Microbiol. Jan 66 (1):23-8.
- Bhugaloo-Vial, P; Douiez, J.P; Moll, D; Dousset, X; Boyaval, P ; Marion, D. (1999). Delineation of key amino acid side chains and peptide domains for antimicrobial properties of Divercin V41, a pediocin-like bacteriocin secreted by *Carnobacterium divergens* V41- Appl. Environ. Microbiol. 65 (7):2895-900.

Bhunia, A.K; Johnson, M.C; Ray, B. and Kalchayanand. (1991). Mode of action of Pediocin AcH from *Pediococcus acidilactici* H on sensitive bacterial strains - J. Appl. Bacteriol. 70: 25-33.

Bradford, M.M. (1976). A rapid and sensitive Method for the Quantitation of Microgram Quantities of Protein Utilizing the principle of Protein-Dye Binding -Anal. Biochem .7 (72): 248-254.

Bruno, M.E.C and Montville, T.J. (1993). Common mechanistic action of bacteriocins from lactic acid bacteria - Appl.Environ.Microbiol. 59: 3003-3010.

Burianek, L.L and Yousef, A.E. (2000). Solvent extraction of bacteriocins from liquid cultures - Lett. Appl. Microbiol . 31(3):193-197.

Chan, W.C; White, P.D. (2000). Fmoc Solid Phase Peptide Synthesis a Practical Approach. Hames B.D (Ed.) Oxford University Press. p: 9-99

Chang, Gu-Gang; Huang, Ter-me; Hung, Hui-Chih. (2000). Reverse Micelles as Life-Mimicking Systems - Proc. Natl. Sci. Counc. Repub. China B. 24 (3): 89-100.

Chatterjee, S; Chatterjee, D.K; Jani, R.H; Blumbach, J; Ganguli, B.N; Klesel, N; Limbert, M; Seibert, G. (1992). Mersacidin, a new antibiotic from *Bacillus*. In vitro and in vivo antibacterial activity – J. Antibiot. (Tokyo). 45:839-845.

Chen, Y; Shapira, R; Eisenstein, M; and Montville, T.J. (1997a). Functional characterization of Pediocin PA-1 binding to liposomes in the absence of a protein receptor and its relation with a predicted tertiary structure - Appl.Environ.Microbiol. 63 (2): 524-531.

Chen, Y; Ludescher, R.D and Montville, T.J. (1997b). Electrostatic interactions, but not the YGNGV consensus motif, govern the binding of Pediocin PA-1 and its fragments to phospholipid vesicles - Appl.Environ.Microbiol. 63 (12): 4770-4777.

Chen, H; and Hoover, D.G. (2003). Bacteriocins and their Food Applications. Comprehensive Reviews in Food Science and Food Safety. 2, 82-99.

Chikindas, M.L; García-Carcera, M.J; Driessens, A.J.M; Ledebotter, A.M; Nissen-Meyer, J; Nes, I.F; Abebe, T; Konings, W.N; and Venema, G. (1993). Pediocin PA-1, a bacteriocin from *Pediococcus acidilactici* PAC1.0, forms hydrophilic pores in the cytoplasmic membrane of target cells - Appl. Environ. Microbiol. 59 (11): 3577-3584.

Chikindas, M; Montville, J. (2002). Perspectives for application of bacteriocins as food preservatives. In: Juneja, V.K Sofos, J.N Editors. Control of foodborne microorganisms. New York: Marcel Dekker Inc.p: 303-21.

Cleveland, J; Montville, T.J; Nes, I.F; Chikindas, M.L (2001). Bacteriocins: Safe, natural antimicrobials for food preservation- Int. J. Food Microbiol. 71: 1-20.

Cordiés jackson, L; Machado Reyes, L.A and Hamilton, M.L. (1998). Principios generales de la terapéutica antimicrobiana- Acta Médica. 8 (1):13-27.

Daeschel, M.A; Mckenney, M.C; McDonald, L.C. (1990). Bactericidal activity of *Lactobacillus plantarum* C-11 - Food. Microbiol. 7: 91-98.

Dathe, M; Nikolenko, H; Meyer, J; Beyermann, M; Bienert, M; (2001). Optimization of the antimicrobial activity of magainin peptides by modification of charge – FEBS.Lett. 501(2-3):146-50.

Dathe, M. and Wieprecht, T. (1999). Structural features of helical antimicrobial peptides: their potential to modulate activity on model membranes and biological cells- Biochim.Biophysica Acta. 1462: 71-87.

DeLano, WL. (2002). The PyMOL Molecular Graphics System DeLano Scientific, San Carlos, CA, USA. <http://www.pymol.org>.

Delves-Broughton, J. (1990). Nisin and its uses as a food preservative - Food Tech. 44: 100-117.

Davey, G.P and Richardson, B.C. (1981). Purification and some properties of diplococin from *Streptococcus cremoris* 346- N .Z.J Dairy Sci. Techno. 16: 84-89.

Dufour, A; Thuault, D; Boulliou, A; Bourgeois, C.M and Le Pennec, J.P. (1991). Plasmid-encoded determinants for bacteriocin production and immunity in a *Lactococcus lactis* strain and purification of the inhibitory peptide- J.Gen. Microbiol. 137 (10):2423-2429.

Dzung, B; and Nes, I.F. (2002). Ribosomally synthesized antibacterial peptides in Gram positive bacteria- Current Drug Targets. 3, 107-122.

Eijsink, V.G.H; Skeie, M; Middelhoven, P.H; Brurberg, M.B; and Nes, I.F. (1998). Comparative studies of Class IIa bacteriocins of lactic acid bacteria- Appl.Environ.Microbiol. 64, 3275-3281.

Eisenberg, D; Weiss, R.M; Terwilliger T.C. (1984). The hydrophobic moment detects periodicity in protein hydrophobicity- Proc. Natl. Acad. Sci. U. S. A. 81,140-144.

Ennahar, S; Sashihara, T; Sonomoto, K. and Ishizaki A. (2000). Class IIa bacteriocins: biosynthesis, structure and activity- FEMS Microbiology Reviews. 24: 85-106.

FDA (U.S. Food & Drug Administration) Federal Register 1988. Nisin preparation: Affirmation of GRAS status as a direct human food ingredient. 21 CFR Part .184, Fed. Reg. 53: 11247-51.

Fields, G.B and Noble, R.I. (1990). Solid phase peptide synthesis utilizing 9-fluorenylmethoxycarbonyl amino acids - Int. J. Pept. Protein Res, Review. 35 (3):161-214.

Fimland, G; Blingsmo, O.R; Sletten, K; Jung, G; Nes, I.F and Nissen-Meyer, J. (1996). New biologically active hybrid bacteriocins constructed by combining regions from various pediocin-like bacteriocins: the C-terminal region is important for determining specificity - Appl. Environ. Microbiol. 62 (9): 3313-3318.

Franz, C.M; Du Toit, M; Olasupo, N.A; Schillinger, U. and Holzapfel, W.H. (1998). Plantaricin D, a bacteriocin produced by *Lactobacillus plantarum* BFE 905 from ready-to-eat-salad- Lett. Appl. Microbiology. 26 (3): 231-235.

Gennis, R.B. (1989). Molecular structure and function- Ed. Springer-Verlag. New York. p.82.

Giangaspero, A ; Sandri, L; Tossi, A. (2001). Amphipathic α -helical antimicrobial peptides- Eur. J. Biochem. 268 (21):5589-5600.

González, C.F and Kunka B.S. (1987). Plasmid-associated bacteriocin production and sucrosed fermentation in *Pediococcus acidilactici*- Appl. Environ. Microbiol. 53:2534-2538.

González, C.F. (1988). Method for inhibiting bacterial spoilage and composition for this purpose- European patent application 88101624.

González, B; Arca, P; Mayo, B; Suarez, J.E. (1994). Detection, purification, and partial characterization of Plantaricin C, a bacteriocin produced by a *Lactobacillus plantarum* strain of dairy origin- Appl. Environ. Microbiol. 60: 2158-2163.

Gratia, A. (1925). Sur un remarquable exemple d'antagonisme entre souches de colibacille- C.R.Soc.Biol. 93:1040-1041.

Hastings, J.W; Sailer, M; Johnson, K; Roy, K.L; Vederas, J.C and Stiles, M.E. (1991). Characterization of Leucocin A-UAL 187 and cloning of the bacteriocin gene from *Leuconostoc gelidum*- J. Bacteriol. 173, 7497-7500.

Hauge, H.H; Mantzilas, D; Moll, G.N; Konings, W.N; Driessen, A.J; Eijsink, V.G; Nissen-Meyer, J. (1998). Plantaricin A is an amphiphilic alpha-helical bacteriocin-like pheromone which exerts antimicrobial and pheromone activities through different mechanisms– Biochemistry. 37 (46):16026-32.

Hauge, H.H; Mantzilas, D; Eijsink, V.G; Nissen-Meyer, J. (1999). Membrane-mimicking entities induce structuring of the two-peptide bacteriocins Plantaricin E/F and Plantaricin J/K – J. Bacteriol. 181 (3):740-747.

Henderson, J.Y; Chopko, A.L; Van Wassenaar, P.D. (1992). Purification and primary structure of Pediocin PA-1 produced by *Pediococcus acidilactici* PAC-10 - Arch. Biochm. Biophys. 295: 5-12.

Hovius, R.; Vallotton, P.; Wohland ,T.; Vogel, H. (2000). Fluorescence techniques: shedding light on ligand-receptor interactions- Trends Pharmacol. Sci. Review. 21(7):266-73.

Holck, A; Axelsson, L; Birkeland, S.E; Aukrust, T; and Blom, H. (1992). Purification and amino acid sequence of Sakasin A, a bacteriocin from *Lactobacillus sake* Lb706- J.Gen.Microbiol. 138, 2715-2720.

Holo, H; Jeknic, Z; Daeschel, M; Stevanovic, S. and Nes, I F. (2001). Plantaricin W from *Lactobacillus plantarum* belongs to a new family of two-peptide lantibiotics - Microbiology. 147: 643-651.

Jacob, F; Lwoff, A; Siminovitch, A. and Wollman, E.L. (1953). Définition de quelques termes relatifs à la lysogénie - Ann.Inst.Pasteur. 84: 222-224.

Jelokhani-Niaraki, M; Prenner, E.J; Kay, C.M; Mc Elhaney, R.N; Hodges, R.S. (2002). Conformation and interaction of the cyclic cationic antimicrobial peptides in lipid bilayers- J.Peptide Res. 60(1): 23-36.

Jiménez-Díaz, R; Piard, J.C; Ruiz-Barba, J.L and Desmazeaud, M.J. (1990). Isolation of a bacteriocin-producing *Lactobacillus plantarum* strain from a green olive fermentation- FEMS.Microbiol.Rev. 87, 91.

Jimenez-Díaz, R; Ríos Sanchez, R.M; Desmazeaud, M; Ruiz Barba, J.L; Piard J.C. (1993). Plantaricin S and T new bacteriocins produced by *Lactobacillus plantarum* LPCO 10 isolated from a green olive fermentation - Appl. Environ. Microbiol. 59: 1416-1424.

Jiménez-Díaz, R.; Maldonado, A; Ruiz-Barba, J.L; (2003). Purification and genetic characterization of Plantaricin NC8, a novel coculture-inducible two-Peptide bacteriocin from *Lactobacillus plantarum* NC8. Appl. Environ. Microbiol. 69 (1): 383-389.

Joerger, M.C and Klaenhammer, T.R (1990). Cloning, expression, and nucleotide sequence of the *Lactobacillus helveticus* 481 gene encoding the bacteriocin Helveticin J. - J.Bacteriol. 172 (11):6339-47.

Joerger, M.C; Hoover, D.G; Barefoot, S.F; Harmon, K.M; Grinstead, D.A; Nettles-Cutter, C.G. (2000). Bacteriocins. In: Ederberg, Editor. Encyclopedia of microbiology, vol 1, 2º Edition. San Diego: Academic Press, Inc. p: 383-97.

Jones, D.T. (1999). Protein Secondary Structure Prediction Based on Position-specific Scoring Matrices- J. Mol. Biol. 292, 195-202.

Jung, G; Sahl, H.G. (1991).Nisin and Novel Lantibiotics. Edited by Jung G., Sahl H.G. Leiden. The Netherlands: ESCOM p: 1-34.

- Kalletta, C; Entian K.D; Jung, G. (1991). Prepeptide sequence of Cinnamycin (Ro09-0198): the first strcuctural gene of a duramycin-type lantibiotic – Eur. J. Biochem. 199 (2):411-415.
- Kato, T; Matsuda, T; Ogawa, E; Ogawa, H; Kato, H; Doi, U. and Nakamura, R. (1994). Plantaricin-149 a Bacteriocin Produced by *Lactobacillus plantarum* NRIC 149 - J. Ferment. Bioeng. 77 (3): 277-282.
- Kelly, S.M. and Price, N.C. (2000). The use of circular dichroism in the investigation of protein structure and function. – Curr. Prot. Pept. Sci. 1 (4): 349-384.
- Kelly, W.J; Asmundson, R.V; Huang, C.M. (1996). Characterization of Plantaricin KW30, a bacteriocin produced by *Lactobacillus plantarum* KW30 – J. Appl. Bacteriol. 81: 657-662.
- Klaenhammer, T.R. (1993). Genetics of bacteriocins produced by lactic acid bacteria - FEMS. Microbiol.Rev.12: 39-86.
- Klaenhammer, T.R (1998). Bacteriocins of lactic acid bacteria- Biochimie. 70, 337-349.
- Konings, W.N; Kok, J; Kuipers, O.P; and Poolman, B. (2000). Lactic acid bacteria: the bugs of the new millennium-Curr.Opin. Microbiol.Review. 3 (3):276-82.
- Konisky, J. (1982). Colicins and other bacteriocins with established mode of action – Annu. Rev. Microbiol. Review. 36: 125-144.
- Kozak, W; Bardowski, J and Dobrzanski, W. (1978). Lactostrepins-acid bacteriocins produced by lactic streptococci -J.Dairy res. 45 (2): 247-257.

Kristiansen, P.E; Fimland, G; Mantzas, D. and Niseen-Meyer, J. (2005). Structure and Mode of Action of the Membrane-permeabilizing Antimicrobial Peptide Pheromone Plantaricin A - J. Biol. Chem. 280 (24): 22945-22950.

Ladanyi, B.M and Faeder, J. (1999). Molecular Dynamics simulations of the interior of aqueous reverse micelles. J.Phys.Chem B. A-N.

Ladokhin, S.A. (2000). Fluorescence Spectroscopy in Peptide and Protein Analysis- Encyclopedia of Analytical Chemistry, R.A Meyers (Ed.) John Wiley & Sons Ltd, Chichester.p: 5762-5779.

Lakowicz, J.R. (1999). Principles of Fluorescence Spectroscopy . Kluwer Academic/New york. pp 1-698.

Laniewska-Moroz, L; Dajnowiec, M; Warminska-Radyko, I. (2001). Effect of *Lactobacillus plantarum* bacteriocin on gram-negative rods - Meded Rijksuniv Gent. Fak Landbouwkd Toegep Biol Wet. 66 (3b):597-600.

Lenz, V.J; Federwisch, M; Gattner, H.G; Bradenburg, D; Höcker, H; Hassiepen,U; Wollmer, A. (1995). Structure and rotational dynamics of fluorescently labeled insulin in aqueous solution and at the amphiphile–water interface of reversed micelles – Biochemistry. 34: 6130-6141.

Lewus, C.B; Sun, S. and Montville, T.J. (1990). Production of an amylase-sensitive bacteriocin by an atypical *Leuconostoc parmesenteroides* strain- Appl.Environ.Microbiol. 58: 143-149.

Lloyd, A.G and Drake, J.J. (1975). Problems posed by essential food preservatives - Br.Med.Bull, Review. 31 (3):214-219.

Lindgren, S.E and Dobrogosz, M.J. (1990). Antagonistic activities of lactic acid bacteria in food and feed fermentations - FEMS Microbiol. Rev. 87: 149-164.

Mackay, L.L; Baldwin, K.A. (1990). Applications for biotechnology: present and future improvements in lactic acid bacteria.-FEMS Microbiol. Rev. Review. 7 (1-2):3-14.

Mayr-Harting, A; Hedges, A.J and Berkley, R.C.W. (1972). Methods for studying bacteriocins - In "Methods in Microbiology" (J.R Norris and D.W Ribbons, Eds.), vol 7^a, p: 315-422. Academic Press, INC., New York.

Merrifield, R.B. (1995). Solid- phase peptide synthesis- In Peptides: Synthesis, Structures, and Applications, Academic Press (San Diego, USA), p: 93-169.

Messi, P; Bondi, M; Sabia, C; Battini, R; Manicardi, G. (2001). Detection and preliminary characterization of a bacteriocin (Plantaricin 35 d) produced by a *Lactobacillus plantarum* strain- Int.J.Food. Microbiol. 64 (1-2): 193-198.

Moll, G; Hildeng-Hauge, H; Nissen-meyer, J; Nes, I.F; Konings, W.N; Driessen, A.J. (1998). Mechanistic Properties of the Two-Component Bacteriocin Lactococcin G - J.Bacteriol. 180 (1): 96-99.

Mor, A. and Nicolas, P.(1994). "The NH₂ terminal α -helical domain 1-18 of Dermaseptin is responsible for antimicrobial activity" - J.Biol. Chem. 269 (3): 1934-1939.

Nissen-Meyer, J; Laesen, A.G; Sletten, K; Daeschel, M; Nes, I.F. (1993). Purification and characterization of Plantaricin A, a *Lactobacillus plantarum* bacteriocin whose activity depends on the action of two peptides - J. Gen. Microbiol. 139 (9): 1973-1978.

Ohmori, N; Niidome, T; Hatakeyama, T; Mihara, H. and Aoyagi, H. (1998). Interaction of α -helical peptides with phospholipid membrane: effects of chain length and hydrophobicity of peptides - J. Pept. Res. 51 p: 103-109.

Ojcius, D.M and Young, J.D.E. (1991). Cytolytic pore-forming proteins and peptides: is there a common structural motif ?- Trends Biochem. Sci.16, 225-229.

Oren, Z; Shai, Y. (1998). Mode of action of linear amphipathic-helical antimicrobial peptides- Biopolymers. 47, 451-453.

Oscáriz, J.C; Pisabarro, A.G. (2001). Classification and mode of action of membrane-active bacteriocins produced by gram-positive bacteria- Int. Microbiol. Review .4:13-19.

Parente, E; Ricciardi, A. (1999). Production, recovery and purification of bacteriocins from lactic acid bacteria.- Appl. Microbiol. Biotechnol. Review. 52 (5): 628-38.

Piard, J.C and Desmazeaud, M.J. (1992). Inhibiting factors produced by lactic acid bacteria, bacteriocins and other antibacterial substances - Lait. 72: 113-142.

Poveda, J.A; Prieto, M; Encinar, J.A; Gonzalez-Ros, J.M; and Reyes Mateo, C. (2003). Intrinsic Tyrosine Fluorescence as a Tool to Study the Interaction of the Shaker B "Ball" Peptide with Anionic Membranes- Biochemistry. 42: 7124-7132.

Quadri, L.E.N; Sailer, M; Roy, K.L; Vederas, J.C and Stiles, M.E. (1994). Chemical and genetic characterization of bacteriocins produced by *Carnobacteriocin piscicola* LV17B- J.Biol.Chem. 269: 12204-12211.

Ray, B. and Daeschel, M. (1992). Food biopreservatives of microbiological origin. CRC Press. Boca Raton.CRC Press.

Reknif, N; Atrihi, A; Lefebvre, G. (1995). Activity of plantaricin SA6, a bacteriocin produced by *Lactobacillus plantarum* SA6 isolated from fermented sausage. - J .Appl. Bacteriol. 78 (4): 349-358.

- Remiger, A; Eijsink, V.G; Ehrmann, M.A; Sletten, K; Nes, I.F; Vogel, RF. (2000). Purification and partial amino acid sequence of plantaricin 1.25 alpha and 1.25 beta, two bacteriocins produced by *Lactobacillus plantarum* TMW 1.25 - J.Appl.Microbiol. 86 (6): 1053-1058.
- Riley, M.A; Wertz, J.E. (2002). Bacteriocin diversity: ecological and evolutionary perspectives - Biochimie. 84 (5-6): 357-364.
- Rodriguez, J.M; Martinez, M.I; Kok, J. (2002). Pediocin PA-1, a wide-spectrum bacteriocin from lactic acid bacteria – Crit. Rev. Food. Sci. Nutr. 42 (2):91-121.
- Sai, K. P; Jagannadham, M. V; Vairamani, M; Raju, N. P. and Sitaram, N. (2001). Tigerinins: novel antimicrobial peptides from the Indian frog Rana Tigerina- J. Biol. Chem. 276 (4): 2701-2707.
- Salminen, S. and Wright, A.V. (1993). Lactic acid bacteria. Future aspects in research and product development on lactic acid bacteria. Ed: Salminen & Wright Marcel Dekker, Inc. New York. p: 429-431.
- Sahl, H.G; Guder, A; Wiedemann, I. (2000). Posttranslationally Modified Bacteriocins The Lantibiotics - Biopolymers (Peptide Science). 55: 62-73.
- Sahl, H.G; Jack, R.W and Biernbaum, G. (1995). Biosynthesis and biological activities of lantibiotics with unique post-translational modifications - Eur. J. Biochem. 230(3): 827-853.
- Shai, Y. (2002). From innate immunity to de-Novo designed antimicrobial peptides - Curr. Pharm. Des. Review. 8(9): 715-725.
- Simonetta, A.C; Carrasco, M.S; Scarinci, H.E. (1997). Inhibitory spectrum of bacteriocin-like substances produced by *Lactobacillus plantarum* strains. - Microbiologie-Aliments-Nutrition. 15: 299-305 .

Sitaram, N. and Nagaraj, R. (1990). A synthetic 13-residue peptide corresponding to the hydrophobic region of bovine seminalplasmin has antibacterial activity and also causes lysis of red blood cells. *J. Biol. Chem.* 18: 10438-10442.

Sobrino, O.J; Rodriguez, J.M; Moreira, W.I; Fernandez, M.F; Sanz, B; Hernandez, P.E. (1991). Antibacterial activity of *Lactobacillus sake* isolated from dry fermented sausages- *Intern. J. Food Microbiol.* 13: 1-10.

Sönnichsen, F.D; Van Eyk, J.E; Hodges, R.S; Sykes, D.B. (1992). Effect of trifluorethanol on protein secondary structure: an NMR and CD study using a synthetic actin peptide- *Biochemistry*. 31 (37): 8790-8798.

Sreerama, N and Woody R. W. (1993). A Self-Consistent Method for the Analysis of Protein Secondary Structure from Circular Dichroism - *Anal. Biochem.* 209 (1): 32-44.

Sreerama, N; Venyaminov, SY; Woody, R.W. (2000b). Estimation of Protein Secondary Structure from Circular Dichroism Spectra: Comparison of CONTIN, SELCON, and CDSSTR methods with an expanded reference set - *Anal. Biochem.* 287: 252–260.

Steinberg, D. A; Hurst, M. A; Fujii, C. A; Kung, A. H; Ho, J. F; Cheng, F. C; Loury, D. J. and Fiddles, J. C. (1997). Protegrin-1: a broad-spectrum, rapidly microbicidal peptide with in vivo activity. *Antimicrob. Agents Chemother.* 41:1738-1742.

Stevens, K.A; Klapes, N.A; Sheldon, B.W; and Klaenhammer, T.R. (1992). Antimicrobial action of nisin against *Salmonella typhimurium* lipopolysaccharide mutants- *Appl. Environ. Microbiol.* 58 (5): 1786-1788

Souto, A.L; Ito, A.S. (2000). Tryptophan fluorescence studies of melanotropins in the amphiphile-water interface of reversed micelles- *Eur. Biophys. J.* 29 (1): 38-47.

Tagg, J.R; and McGiven, A.R. (1971). Assay system for bacteriocins- *Appl. Environ. Microbiol.* 21: 943-947.

Tagg, J.R; Dajani, A.S and Wannamaker, L.W. (1976). Bacteriocins of Gram-positive bacteria - Bacteriol.Rev. 40 (3): 722-756.

Tagg, J.R. (1991). Bacterial BLIS. ASM News. 57:611.

Talarico, T.L; Casas, I.A; Chung, T.C; Dobrogosz, W.J. (1988). Production and isolation of reuterin, a growth inhibitor produced by *Lactobacillus reuteri*—Antimicrob. Agents Chemother. 32: 1854-1858.

Todorov, S; Onno, B; Sorokine, O; Chobert, J.M; Ivanova, I; Dousset, X. (1999). Detection and characterization of a novel antibacterial substance produced by *Lactobacillus plantarum* ST31 isolated from sourdough- Int.J.Food.Microbiol. 48(3): 167-177.

Todorov, S.D; Dicks, L.M. (2005). Characterization of bacteriocins produced by lactic acid bacteria isolated from spoiled black olives– J.Basic Microbiol. 45 (4):312-22.

Turner, D.L; Brennan, L; Meyer, H.E; Lohaus, C; Siethoff, C; Costa, H.S; Gonzalez, B; Santos, H; Suarez, J.E. (1999). Solution structure of Plantaricin C, a novel lantibiotic- Eur. J. Biochem. 264 (3): 833-839.

Uteng, M; Hauge, H.H; Markwick, P; Fimland, G; Mantzilas, D; Nissen Meyer, J; Muhle-Goll, C. (2003). Three dimensional structure in lipid micelles of the Pediocin-Like antimicrobial peptide Sakacin P- Biochemistry. 42, 11417 (Primary citation). On Line: <http://www.rcsb.org/pdb/explore.do?structureId=10G7>.

Varadaraj, M.C; Suma, K; Misra, M.C. (1998). Plantaricin LP84, a broad spectrum heat-stable bacteriocin of *Lactobacillus plantarum* NCIM 2084 produced in simple glucose broth medium- Int.J. Food Microbiol. 40 (1-2): 17-25.

- Van Reenen, C.A; Dicks, L.M; Chikindas, M.L. (1998). Isolation, purification and partial characterization of Plantaricin 423, a bacteriocin produced by *Lactobacillus plantarum*- J.Appl.Microbiol.4(6):1131-7.
- Vandenbergh, P.A; Pucci, M.J; Kunka, B.S and Vedamuthu, E.R. (1989). Method for inhibiting *Listeria monocytogenes* using a bacteriocin- European patent application 89101125.6.
- Vignolo, G.M; Suriani, F; Pesce de Ruiz Holgado, A; Oliver, G. (1993). Antibacterial activity of *Lactobacillus plantarum* strains isolated from dry fermented sausages - J. Appl. Bacteriol. 75 (4): 344-349.
- Yang, R; Johnson, M.C. and Ray, B. (1992). Novel method to extract large amounts of bacteriocins from lactic acid bacteria- Appl. Environ. Microbiol. 58 (10): 3355-3359.
- Wade, J.D; Bedford, J; Sheppard, R.C; Tregear, G.W. (1991). DBU as an N-alpha-deprotecting reagent for the fluorenylmethoxycarbonyl group in continuous flow solid-phase peptide synthesis.- Pept Res. (3):194-9.
- West, C.A and Warner, P.J. (1988). Plantaricin B, a bacteriocin produced by *Lactobacillus plantarum* NCDO 1193- FEMS. Microbiol. Lett. 49: 163-165.
- Woody, R. W. (1996). Theory of circular dichroism of proteins. In Fasman G.D. (ed). Dichroism and the Conformational Analysis of Biomolecules - Plenum Press, New York/London. p: 25-67.
- Wu, M. and Hancock, R.E.W. (1999) Interaction of the cyclic antimicrobial cationic peptide bactenecin with the outer and cytoplasmic membrane. J. Biol. Chem. 274: 29-35.

- Zhang, L (2001). Innovation & Perspectives in Solid Phase Synthesis, 3rd International Symposium”, R. Epton (ed.) Mayflower Scientific Ltd., Birmingham, pp. 711.
- Zhou, H.; Zhou Y. (2005). Fold recognition by combining sequence profiles derived from evolution and from depth-dependent structural alignment of fragments. Proteins. 58: 321-328.